



HAL
open science

Gender Differences in the Association between Socioeconomic Status and Subclinical Atherosclerosis

Olivier Grimaud, Annabelle Lapostolle, Claudine Berr, Catherine Helmer, Carole Dufouil, Wahida Kihal, Annick Alperovitch, Pierre Chauvin

► **To cite this version:**

Olivier Grimaud, Annabelle Lapostolle, Claudine Berr, Catherine Helmer, Carole Dufouil, et al.. Gender Differences in the Association between Socioeconomic Status and Subclinical Atherosclerosis. PLoS ONE, 2013, 8 (11), pp.e80195. 10.1371/journal.pone.0080195 . hal-02437437

HAL Id: hal-02437437

<https://ehesp.hal.science/hal-02437437>

Submitted on 26 May 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Gender Differences in the Association between Socioeconomic Status and Subclinical Atherosclerosis

Olivier Grimaud^{1*}, Annabelle Lapostolle², Claudine Berr³, Catherine Helmer⁴, Carole Dufouil^{5,6}, Wahida Kihal¹, Annick Alperovitch^{5,6}, Pierre Chauvin²

1 French School of Public Health, Rennes, France, **2** Institut National de la Santé et de la Recherche Médicale U707, Paris, France, **3** Institut National de la Santé et de la Recherche Médicale, U1061, Montpellier, France, **4** Institut National de la Santé et de la Recherche Médicale, U897, Bordeaux, France, **5** Université Bordeaux Segalen, Bordeaux, France, **6** Institut National de la Santé et de la Recherche Médicale 708, Paris, France

Abstract

Objectives: This study explored the pattern of associations between socioeconomic status (SES) and atherosclerosis progression (as indicated by carotid intima media thickness, CIMT) across gender.

Design: Cross-sectional analysis of a sample of 5474 older persons (mean age 73 years) recruited between 1999 and 2001 in the 3C study (France). We fitted linear regression models including neighborhood SES, individual SES and cardiovascular risk factors.

Results: CIMT was on average 24 μm higher in men (95% CI: 17 to 31). Neighborhood SES was inversely associated with CIMT in women only (highest versus lowest tertiles: $-12.2 \mu\text{m}$, 95%CI -22 to -2.4). This association persisted when individual SES and risk factors were accounted for. High individual education was associated with lower CIMT in men ($-21.4 \mu\text{m}$ 95%CI -37.5 to -5.3) whereas high professional status was linked to lower CIMT among women ($-15.7 \mu\text{m}$ 95%CI: -29.2 to -2.2). Adjustment for cardiovascular risk factors resulted in a slightly more pronounced reduction of the individual SES-CIMT association observed in men than in women.

Conclusion: In this sample, neighborhood and individual SES displayed different patterns of associations with subclinical atherosclerosis across gender. This suggests that the causal pathways leading to SES variations in atherosclerosis may differ among men and women.

Citation: Grimaud O, Lapostolle A, Berr C, Helmer C, Dufouil C, et al. (2013) Gender Differences in the Association between Socioeconomic Status and Subclinical Atherosclerosis. PLoS ONE 8(11): e80195. doi:10.1371/journal.pone.0080195

Editor: Angelo Scuteri, INRCA, Italy

Received: June 19, 2013; **Accepted:** September 30, 2013; **Published:** November 25, 2013

Copyright: © 2013 Grimaud et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The Three-City Study is conducted under a partnership agreement between the Institut National de la Santé et de la Recherche Médicale (INSERM), the Victor Segalen – Bordeaux II University, and Sanofi-Aventis. The Fondation pour la Recherche Médicale funded the preparation and initiation of the study. The 3C Study is also supported by the Caisse Nationale Maladie des Travailleurs Salariés, Direction Générale de la Santé, MGEN, Institut de la Longévité, Conseils Régionaux of Aquitaine and Bourgogne, Fondation de France, and Ministry of Research–INSERM Programme “Cohortes et collections de données biologiques.” The funding organizations played no role in the design and conduct of the study; in the collection, management, analysis, and interpretation of the data; or in preparation, review, or approval of the manuscript.

Competing Interests: The 3C study has received funding from various sources including a commercial source (Sanofi-Aventis). This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: olivier.grimaud@ehesp.fr

Introduction

There is good evidence that individual socioeconomic status (SES) is associated with cardiovascular risk. The gradient of increasing risk when SES decreases has long been demonstrated in studies measuring cardiovascular events such as myocardial infarction [1] and stroke [2]. There is also evidence that contextual factors, for instance the SES of the area of residence, are associated with cardiovascular health, which is worsened in socially deprived neighborhoods after controlling for individual SES [3]. More recently, studies have explored the link between SES and subclinical atherosclerosis as measured by the carotid intima media thickness (CIMT). CIMT measurement by ultrasound is relatively simple, non-invasive and correlates well with histology [4]. In their meta-analysis, Lorenz et al showed that a 0.10 mm increase of the common carotid intima media thickness

corresponds to a 1.15 relative risk of developing myocardial infarction (95% CI, 1.12 to 1.17), and 1.18 relative risk for stroke (95% CI 1.16 to 1.21) [5]. Previous studies have shown that low education, low income, or manual occupation are associated with thicker carotid arterial wall [6–11] or with faster progression of carotid wall thickness [12]. Neighborhood SES measured either instantaneously [13–16] or over the lifecourse [17,18] also shows an inverse association with CIMT.

To date the few studies which have investigated the link between SES and CIMT have been conducted in the USA and in Northern Europe. Since evidence in this domain of research only rely on observational studies, findings from different epidemiological settings help strengthening the case for a causal link. Our first aim was therefore to examine whether similar associations existed in France, a country where cardiovascular diseases (CVD) represent a less prominent part of the disease burden, [19] but is

not void of significant health inequalities [20]. Cardiovascular diseases develop 7 to 10 years later in women, most probably as a consequence of exposure to endogenous estrogens and lower prevalence of risk factors [21]. Since results from earlier studies suggested that the influence of SES on the development of subclinical atherosclerosis differs in men and women [11,14,17] we also examined the pattern of the SES-CIMT associations across gender.

Methods

The 3C study is an ongoing longitudinal study on vascular risk factors and dementia. For the present cross sectional analysis we used information collected at baseline. Full description of the study design has been reported elsewhere [22]. Briefly, between 1999 and 2001, non-institutionalized persons aged 65 years and over were recruited from the electoral rolls of Bordeaux, Dijon and Montpellier. Eligible inhabitants were invited to participate and the overall acceptance rate was 37% giving a total number of 9294. Each participant signed an informed consent. The study protocol received approval from the Ethical Committee of the University Hospital of Kremlin-Bicêtre. Ultrasound examination of the carotid arteries was proposed only to participants less than 85 who were able to come to the examination centers. For costs constraints, measurements could not be performed in the last 4 months of the baseline phase of the study. Overall, valid ultrasound measurements were available for 69.5% of the participants ($n = 6462$).

Measurement of carotid intima media thickness

As described elsewhere [23,24] ultrasound examinations followed a common protocol and all results were read centrally at the reference center (Hôpital Broussais, Paris). Each study center used a B-mode system (Ultramark 9 High Definition Imaging) with a 5 to 10-MHz sounding. Carotid intima media thickness (CIMT) was measured at a plaque free site along a 10-mm-long segment of the far wall of right and left common carotid arteries 2 to 3 cm proximal to the bifurcation. It was measured as the distance between the lumen-intima interface and the media-adventitia interface using an automated edge detection algorithm. On average, 75 measurements were automatically performed on each side. The CIMT used in the analysis corresponds to the mean of the right and left mean values. A reproducibility study was conducted on 114 subjects who underwent 2 ultrasound examinations performed blindly by 2 different sonographers. The results showed a mean absolute difference of 60 μm and a correlation coefficient of 0.71 between repeated examinations which is comparable to figures found in other large community studies [25,26].

Individual and neighborhood socioeconomic variables

Education level was collected in 6 categories ranging from no diploma to university degree that we grouped in 3 classes: less than 6 years, 6 to 11 years and ≥ 12 years of schooling. Professional status was also grouped in 3 categories: high (managers, professionals, technicians and associate professionals) intermediate (clerical, service, shop and market sale workers, armed forces) and low (all other professions including housewife). Household income was initially collected in 4 categories. Since only 272 participants (5%) declared monthly income of less than 750€ per month, these were grouped to the next level of income in order to obtain 3 groups of comparable size (<1500€, 1500-2250€, >2250€).

Postal addresses were geocoded in order to match participants to their neighborhood of residence. As done in other studies

[27,28], we used IRIS (for “*Ilots Regroupés pour l’Information Statistiques*”) as proxy for neighborhoods. IRIS is the smallest census aggregation level used by the French national statistical office to disseminate information. It corresponds to homogenous geographical areas with borders that follow main roads or other physical features such as rivers or railways. At the time of the 1999 population census in France, the average population size of IRISs was 2400. Because Bordeaux, Dijon and Montpellier represent socially and culturally contrasted urban settings and displayed significantly different values of mean CIMT, we tried to identify a common neighborhood SES indicators which fulfilled the two following properties: 1) showed a large range of values between IRISs in each center so as to increase discrimination power, 2) displayed comparable distributions across centers (as reflected by medians and interquartile values) so as to limit confounding. We used the 1999 census and the 2001 “household tax income” data files provided by the French National Office of Statistics to derive unemployment rate, proportion of manual workers, median annual income, proportion of adults with secondary education and the Townsend deprivation score. The proportion of adults with secondary education proved to best fit our two criteria and was therefore selected as neighborhood SES indicator.

Risk factors variables

Measurements of height, weight, systolic and diastolic blood pressure, were performed during clinical examinations. One laboratory carried out measurements of all biological parameters, including fasting glycaemia and lipid levels. We entered diabetes as a 3 categories variable (glycaemia <6.1 mmol/l; $6.1 \leq$ glycaemia ≤ 7 mmol/l; glycaemia >7 mmol/l or treatment), hypercholesterolemia as a binary variable (serum total cholesterol > 6.2 mmol/l, or treatment), systolic blood pressure in five categories (quintiles) and added a variable indicating use of antihypertensive treatment. Smoking status was included as former, current or none, alcohol consumption in 4 categories (abstinent, <10, <20 or ≥ 20 gr/day), and BMI (weight/height²) as a continuous variable.

Statistical analysis

We first examined the distribution of individual SES variables, risks factors and mean CIMT. We used the Chi² test and the Kruskal Wallis test to determine statistical significance of the differences across gender. We analyzed men and women separately and tested for gender SES interactions in models including all participants when results suggested differing associations across genders. Since multi-level models showed little evidence of clustering (intraclass correlation lower than 1%) we estimated parameters using single level multiple linear regression models. All models were adjusted for age and center. We started by exploring associations between CIMT and neighborhood SES categorized in tertiles. We then added individual SES variables in order to estimate their own associations with CIMT and their influence on the neighborhood SES-CIMT relationship. To better ascertain the influence of income, we fitted models restricted on the subsample of participants living on their own for whom household income therefore does not depend of household composition. In the final stage of the analysis we checked whether the associations between SES and CIMT were modified when accounting for cardiovascular risk factors. All analyses were performed with Stata v10.2 (StataCorp LP, TX, USA).

Table 1. Means and percent distributions of socioeconomic status indicators and cardiovascular risk factors for the entire sample and by gender.

	Entire sample	Men	Women	p*
Number of participants	5 474	2 007	3 467	
Age (mean (sd) in years)	73.3 (4.9)	73.2 (4.9)	73.4 (4.8)	0.30
Living alone (%)	35.5	13.5	48.3	<10 ⁻³
Participant socioeconomic status (%)				
Education				<10 ⁻³
<6 years	31.2	26.8	33.8	
6–11 years	30.3	26.5	32.5	
≥12 years of schooling	38.5	46.7	33.7	
Professional status				<10 ⁻³
Low	38.8	32.6	42.4	
Intermediate	40.6	31.9	45.6	
High	20.6	35.4	12.0	
Household monthly income				<10 ⁻³
Low (<1 500€)	34.9	19.4	44.1	
Medium (1 500 to 2250€)	29.3	31.1	28.3	
High (>2 250€)	35.8	49.5	27.6	
Neighborhood socioeconomic status				
Number of neighborhoods	242			
Proportion of adults with secondary education in the neighborhoods (mean (sd))	53 (15)			
Behavioral risk factors				
Smoking (% former or current)	36.7	67.8	18.8	<10 ⁻³
Mean (sd) alcohol consumption (gr/day)	12.9 (14.7)	21.8 (18.2)	7.8 (8.9)	<10 ⁻³
Mean (sd) body mass index (kg/m ²)	25.5 (4.0)	26.1 (3.4)	25.1 (4.2)	<10 ⁻³
Biological risk factors				
Mean (sd)systolic blood pressure (mm Hg)	145 (21)	150 (21)	142 (21)	<10 ⁻³
Receiving Anti-hypertensive therapy (%)	43.5	42.1	44.3	0.11
Diabetes (%)	8.5	11.8	6.6	<10 ⁻³
Hypercholesterolaemia (%)	55.0	41.7	62.7	<10 ⁻³
Carotid intima-media thickness				
Mean (sd) in μm	711 (120)	727 (129)	702 (113)	<10 ⁻³

3C study (France, 1999–2001).

*p-value for differences between gender (Kruskal-Wallis test for continuous variables and Chi² test for categorical variables).

doi:10.1371/journal.pone.0080195.t001

Results

Valid ultrasound measures were available for 6462 participants. We excluded those having a self-reported history of cardiovascular disease ($n = 847$) and those who could not be matched to their neighborhood of residence ($n = 110$). Because estimation of the variance at the neighborhood level requires the presence of at least two participants in a given neighborhood, a further 31 “isolated” records were excluded. The final study sample consists thus of 5474 individuals (2007 men and 3467 women) with mean age 73 years, distributed in 242 neighborhoods. Mean CIMT was highest in Bordeaux (749 μm), intermediate in Montpellier (712 μm) and lowest in Dijon (695 μm, $p < 10^{-3}$).

Information on individual SES was missing for 324 (5.9%) participants, mostly because of missing household income ($n = 305$), who therefore could not be included in some models. Participants with missing individual SES were more often women (77% versus 62% among others, $p < 10^{-3}$) and older (mean difference +1.6 year, $p < 10^{-3}$) but had similar CIMT.

Men and women differed markedly according to socioeconomic status and risk factors profile (table 1). Men were more educated and had higher professional status. Nearly twice as many men than women declared household income in the highest category but this should be seen in light of the fact that women were more often living on their own (48.3 versus 13.5%). Smoking and alcohol drinking were much more frequent among men and overall, with the exception of hypercholesterolemia, their biological risk factors profile was less favorable. Men’s carotid wall was on average 24 μm (95% CI: 17 to 31) thicker than that of women.

Neighborhood SES showed a significant association with CIMT in women (table 2). Among them, each increase in the tertile of neighborhood education corresponded to a decrease of approximately 6 μm of CIMT (linear trend in model 1, $p = 0.02$). This association persisted when individual SES (model 2) and risk factors (model 3) were accounted for. In men, although CIMT decreased with increasing level of neighborhood in model 1, this failed to reach statistical significance (linear trend, $p = 0.30$).

Table 2. Associations between socioeconomic status and carotid intima media thickness. Values are differences in carotid intima media thickness measured in μm (95% CI) 3C study (France, 1999–2001).

	Men			Women		
	Model 1 neighborhood SES (N = 2007)	Model 2 model 1 + individual SES (N = 1933)	Model 3 model 2 + risk factors ^a (N = 1933)	Model 1 neighborhood SES (N = 3467)	Model 2 model 1 + individual SES (N = 3217)	Model 3 model 2 + risk factors ^a (N = 3217)
Neighborhood SES ^b						
1st tertile (reference)	0	0	0	0	0	0
2nd tertile	−4.5 (−18.0; 9.0)	−5.8 (−19.8; 8.1)	−2.9 (−16.8; 10.9)	−7.1 (−16.1; 1.8)	−5.7 (−15.0; 3.9)	−4.5 (−13.8; 5.0)
3rd tertile	−7.6 (−22.1; 6.8)	−5.9 (−21.2; 9.5)	−3.5 (−18.7; 11.7)	−12.2 * (−22.0; −2.4)	−12.4 * (−22.8; −2.0)	−10.8 * (−21.1; −0.4)
Individual education						
< 6 years (reference)		0	0		0	0
6–11 years		1.4 (−14.1; 17.0)	5.1 (−10.4; 20.5)		−7.9 (−17.7; 1.8)	−6.8 (−16.6; 2.9)
≥12 years		−21.4 ** (−37.5; −5.3)	−16.9* (−32.9; −0.9)		−2.9 (−13.7; 7.9)	−0.7 (−11.6; 10.1)
Professional status						
Low (reference)		0	0		0	0
Intermediate		−0.3 (−14.7; 14.1)	3 (−11.3; 17.2)		−10.8* (−19.4; −2.1)	−9.8* (−18.3; −1.2)
High		9.3 (−6.7; 25.3)	9.6 (−6.3; 25.4)		−15.7* (−29.2; −2.2)	−13.3 (−26.8; 0.1)
Household income						
Low (reference)		0	0		0	0
Medium		3.2 (−13.1; 19.6)	2.5 (−13.7; 18.7)		−0.7 (−10.0; 8.6)	1.3 (−8.0; 10.6)
High		12.2 (−4.6; 29.1)	12.2 (−4.5; 28.9)		3.8 (−6.4; 14.0)	6.2 (−4.0; 16.4)

* $p < 0.05$ ** $p < 0.01$.

All models are adjusted for age and center.

^asmoking; alcohol intake; body mass index; systolic BP; antihypertensive treatment; diabetes; hypercholesterolemia.^bSocioeconomic status of the neighborhood measured as the proportion of adults with secondary education.

doi:10.1371/journal.pone.0080195.t002

Adjustment for individual SES and risk factors tended to attenuate the association. No significant gender-neighborhood SES interaction existed in model including both men and women.

The pattern of individual SES and CIMT associations also varied across gender. In men, no difference in CIMT was noticeable between the first two levels of education. However, compared to those with less than 6 years of schooling, those with 12 years or more had lower CIMT ($-21.4 \mu\text{m}$, $p = 0.009$). Adjustment for risk factors (model 3) reduced this association by 21%. Results regarding individual income in men were suggestive of a positive association with CIMT but this association failed to reach statistical significance (p for linear trend = 0.13).

In contrast with men, only professional status was associated with CIMT among women (gender-professional status interaction: $p < 0.05$). CIMT decreased as professional status increased and was significantly lower among women who had a non-manual profession (p for linear trend 0.007 in model 2 and 0.02 in model 3). This relationship was only slightly reduced when adjusting for risk factors. Only 498 women of our sample (14%) declared to be

housewife and excluding them from the analysis tended to accentuate the association between non manual profession and CIMT (not shown). There was no discernible association between household income and CIMT, whether all women (model 2 and 3) or only those living alone were analyzed (not shown). Too few men were living on their own ($n = 271$) for us to derive a precise estimate of the income CIMT association in this subgroup.

Discussion

We showed that the inverse association between socioeconomic status and subclinical atherosclerosis observed in the USA and northern Europe also applies to France, a country where cardiovascular diseases are less prevalent. Similar to others, we found that both individual and contextual SES factors contribute to this relationship. An interesting aspect of our results is the differential pattern of these associations across gender.

First, neighborhood SES showed a clearer negative association with CIMT in women than in men and this association was statistically significant only in the former. It is possible that our

sample lacks statistical power in order to identify significant association in men. However, previous studies have shown a similar stronger effect among women of neighborhood SES on subclinical atherosclerosis [14,17], incidence of coronary heart diseases [29] and self-rated health [30]. In adult life, women's involvement in childcare and part-time work is likely to have resulted in them being more exposed than men to the physical and social aspects of their area of residence. Men and women may also differ in the ways they perceive their local environment and make use of health promoting services and facilities [30]. Neighborhoods that we classified as advantaged in our study on the basis of residents' educational level are likely to offer more or higher quality of these services and facilities.

Our results also showed differences across gender with respect to the individual SES indicator associated with CIMT, namely education in men and professional status in women. Moreover, adjustment for cardiovascular risk factors suggests that they may explain part of the education-CIMT association in men, but less so of the profession-CIMT association in women. Evidence is accumulating regarding the role of psychosocial factors in the development of ischemic heart diseases [31]. For instance, work and marital stress were positively associated with faster progression of atherosclerosis in a prospective study of women hospitalized for an acute coronary event [32]. Results from animal studies indicate that the relation between social status and CVD may be sex-specific [33]. Specifically, social subordination may increase the risk of atherosclerosis in female but not in male monkeys, which somehow corroborates our findings regarding occupation. More generally, women may be more exposed and perhaps more susceptible to psychological stressors from their neighborhood, family and professional environments, whereas the contribution of adverse health behaviors might be more important in men of this generation. Taken together these results are compatible with the hypothesis that the causal pathways leading to social disparities in atherosclerosis differ across gender.

Strengths and limitations

Strengths of our study include the large sample size, standardized measurements for CIMT and other co-variables, and

availability of both individual and neighborhood SES. Assessment of the latter may suffer from measurement error however, since lifetime (as opposed to current) exposure to neighborhood deprivation is more closely linked to subclinical atherosclerosis [14,17]. Also, we had no information on non-residential neighborhood exposures which have been shown to confound the association between health and residential neighborhood [34]. Our results may therefore underestimate the strength of neighborhood CIMT associations. Similarly, relying on current as opposed to lifetime exposure to risk factors may underestimate their contributions to social health inequalities [35]. The 3C study's initial 37% acceptance rate raises the issue of selection bias, which would arise if acceptance was differentially linked to health across SES [36]. Assuming that higher SES individuals were more willing to take part to 3C despite lower health level, our results would again underestimate the "true" negative association between SES and CIMT. Finally, because of the cross-sectional design of our study, the associations we identified do not necessarily reflect causal relationships.

Conclusion

In conclusion, this study adds to the body of evidence suggesting that both contextual and individual SES play a part in the development of subclinical atherosclerosis. The pattern of gender differences that we observed is consistent with the hypothesis that, at the time of data collection, the causal pathways leading to social inequalities in atherosclerosis differed across gender. Although a different mix of exposure to social and cardiovascular risk factors may now prevail, public health actions aiming at reducing these inequalities may still need adaptation to these differences.

Author Contributions

Conceived and designed the experiments: OG PC. Analyzed the data: OG AL WK. Wrote the paper: OG. As initiators of the 3C study, supervised the analysis examined and discussed interim results, commented on earlier drafts and approved the final version of the paper: CB CD AA CH.

References

- Gonzalez MA, Rodriguez AF, Calero JR (1998) Relationship between socioeconomic status and ischaemic heart disease in cohort and case-control studies: 1960–1993. *Int J Epidemiol* 27: 350–358.
- Cox AM, McKeivitt C, Rudd AG, Wolfe CD (2006) Socioeconomic status and stroke. *Lancet Neurol* 5: 181–188.
- Chaix B (2009) Geographic life environments and coronary heart disease: a literature review, theoretical contributions, methodological updates, and a research agenda. *Annu Rev Public Health* 30: 81–105.
- Pignoli P, Tremoli E, Poli A, Oreste P, Paoletti R (1986) Intimal plus medial thickness of the arterial wall: a direct measurement with ultrasound imaging. *Circulation* 74: 1399–1406.
- Lorenz MW, Markus HS, Bots ML, Rosvall M, Sitzer M (2007) Prediction of clinical cardiovascular events with carotid intima-media thickness: a systematic review and meta-analysis. *Circulation* 115: 459–467.
- Diez-Roux AV, Nieto FJ, Tyroler HA, Crum LD, Szklo M (1995) Social inequalities and atherosclerosis. The atherosclerosis risk in communities study. *Am J Epidemiol* 141: 960–972.
- Lynch J, Kaplan GA, Salonen R, Cohen RD, Salonen JT (1995) Socioeconomic status and carotid atherosclerosis. *Circulation* 92: 1786–1792.
- Ebrahim S, Papacosta O, Whincup P, Wannamethee G, Walker M et al. (1999) Carotid plaque, intima media thickness, cardiovascular risk factors, and prevalent cardiovascular disease in men and women: the British Regional Heart Study. *Stroke* 30: 841–850.
- Lamont D, Parker L, White M, Unwin N, Bennett SM et al. (2000) Risk of cardiovascular disease measured by carotid intima-media thickness at age 49–51: lifecourse study. *BMJ* 320: 273–278.
- Rosvall M, Ostergren PO, Hedblad B, Isacson SO, Janzon L et al. (2000) Occupational status, educational level, and the prevalence of carotid atherosclerosis in a general population sample of middle-aged Swedish men and women: results from the Malmö Diet and Cancer Study. *Am J Epidemiol* 152: 334–346.
- Nash SD, Cruickshanks KJ, Klein R, Klein BE, Nieto FJ et al. (2011) Socioeconomic status and subclinical atherosclerosis in older adults. *Prev Med* 52: 208–212.
- Lynch J, Kaplan GA, Salonen R, Salonen JT (1997) Socioeconomic status and progression of carotid atherosclerosis. Prospective evidence from the Kuopio Ischemic Heart Disease Risk Factor Study. *Arterioscler Thromb Vasc Biol* 17: 513–519.
- Deans KA, Bezlyak V, Ford I, Batty GD, Burns H et al. (2009) Differences in atherosclerosis according to area level socioeconomic deprivation: cross sectional, population based study. *BMJ* 339: b4170.
- Lemelin ET, Diez Roux AV, Franklin TG, Carnethon M, Lutsey PL et al. (2009) Life-course socioeconomic positions and subclinical atherosclerosis in the multi-ethnic study of atherosclerosis. *Soc Sci Med* 68: 444–451.
- Ranjit N, Diez-Roux AV, Chambless L, Jacobs DR Jr, Nieto FJ et al. (2006) Socioeconomic differences in progression of carotid intima-media thickness in the Atherosclerosis Risk in Communities study. *Arterioscler Thromb Vasc Biol* 26: 411–416.
- Nordstrom CK, Diez Roux AV, Jackson SA, Gardin JM (2004) The association of personal and neighborhood socioeconomic indicators with subclinical cardiovascular disease in an elderly cohort. The cardiovascular health study. *Soc Sci Med* 59: 2139–2147.
- Murray ET, Diez Roux AV, Carnethon M, Lutsey PL, Ni H et al. (2010) Trajectories of neighborhood poverty and associations with subclinical atherosclerosis and associated risk factors: the multi-ethnic study of atherosclerosis. *Am J Epidemiol* 171: 1099–1108.
- Carson AP, Rose KM, Catellier DJ, Kaufman JS, Wyatt SB et al. (2007) Cumulative socioeconomic status across the life course and subclinical atherosclerosis. *Ann Epidemiol* 17: 296–303.

19. Sutton CJ, Marsden J, Watkins CL, Leathley MJ, Dey P (2010) Changing stroke mortality trends in middle-aged people: an age-period-cohort analysis of routine mortality data in persons aged 40 to 69 in England. *J Epidemiol Community Health* 64: 523–529.
20. Mackenbach JP, Stirbu I, Roskam AJ, Schaap MM, Menvielle G et al. (2008) Socioeconomic inequalities in health in 22 European countries. *N Engl J Med* 358: 2468–2481.
21. Maas AH, Appelman YE (2010) Gender differences in coronary heart disease. *Neth Heart J* 18: 598–602.
22. The 3C Study Group (2003) Vascular factors and risk of dementia: design of the Three-City Study and baseline characteristics of the study population. *Neuroepidemiology* 22: 316–325.
23. Zureik M, Gariepy J, Courbon D, Dartigues JF, Ritchie K et al. (2004) Alcohol consumption and carotid artery structure in older French adults: the Three-City Study. *Stroke* 35: 2770–2775.
24. Plichart M, Celermajer DS, Zureik M, Helmer C, Jouven X et al. (2011) Carotid intima-media thickness in plaque-free site, carotid plaques and coronary heart disease risk prediction in older adults. The Three-City Study. *Atherosclerosis* 219: 917–924.
25. O'Leary DH, Polak JF, Wolfson SK Jr, Bond MG, Bommer W et al. (1991) Use of sonography to evaluate carotid atherosclerosis in the elderly. The Cardiovascular Health Study. CHS Collaborative Research Group. *Stroke* 22: 1155–1163.
26. Bots ML, Mulder PG, Hofman A, van Es GA, Grobbee DE (1994) Reproducibility of carotid vessel wall thickness measurements. The Rotterdam Study. *J Clin Epidemiol* 47: 921–930.
27. Deguen S, Lalloue B, Bard D, Havard S, Arveiler D et al. (2010) A small-area ecologic study of myocardial infarction, neighborhood deprivation, and sex: a Bayesian modeling approach. *Epidemiology* 21: 459–466.
28. Grimaud O, Bejot Y, Heritage Z, Vallec J, Durier J et al. (2011) Incidence of stroke and socioeconomic neighborhood characteristics: an ecological analysis of dijon stroke registry. *Stroke* 42: 1201–1206.
29. Sundquist K, Malmstrom M, Johansson SE (2004) Neighbourhood deprivation and incidence of coronary heart disease: a multilevel study of 2.6 million women and men in Sweden. *J Epidemiol Community Health* 58: 71–77.
30. Stafford M, Cummins S, Macintyre S, Ellaway A, Marmot M (2005) Gender differences in the associations between health and neighbourhood environment. *Soc Sci Med* 60: 1681–1692.
31. Vaccarino V, Badimon L, Corti R, de Wit C, Dorobantu M et al. (2011) Ischaemic heart disease in women: are there sex differences in pathophysiology and risk factors? Position paper from the working group on coronary pathophysiology and microcirculation of the European Society of Cardiology. *Cardiovasc Res* 90: 9–17.
32. Wang HX, Leineweber C, Kirkeide R, Svane B, Schenck-Gustafsson K et al. (2007) Psychosocial stress and atherosclerosis: family and work stress accelerate progression of coronary disease in women. The Stockholm Female Coronary Angiography Study. *J Intern Med* 261: 245–254.
33. Petticrew M, Davey SG (2012) The monkey puzzle: a systematic review of studies of stress, social hierarchies, and heart disease in monkeys. *PLoS One* 7: e27939.
34. Inagami S, Cohen DA, Finch BK (2007) Non-residential neighborhood exposures suppress neighborhood effects on self-rated health. *Soc Sci Med* 65: 1779–1791.
35. Stringhini S, Sabia S, Shipley M, Brunner E, Nabi H et al. (2010) Association of socioeconomic position with health behaviors and mortality. *JAMA* 303: 1159–1166.
36. Larsen SB, Dalton SO, Schuz J, Christensen J, Overvad K et al. (2012) Mortality among participants and non-participants in a prospective cohort study. *Eur J Epidemiol* 27: 837–845.