

Demographic and socio-economic factors related to food intake and adherence to nutritional recommendations in a cohort of pre-school children

Lise Dubois, Anna Farmer, Manon Girard, Daniel Burnier, Marion Porcherie

► **To cite this version:**

Lise Dubois, Anna Farmer, Manon Girard, Daniel Burnier, Marion Porcherie. Demographic and socio-economic factors related to food intake and adherence to nutritional recommendations in a cohort of pre-school children. Public Health Nutrition, Cambridge University Press (CUP), 2011, 14 (6), pp.1096-1104. 10.1017/S1368980010003769 . hal-02884260

HAL Id: hal-02884260

<https://hal.ehesp.fr/hal-02884260>

Submitted on 30 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Demographic and socio-economic factors related to food intake and adherence to nutritional recommendations in a cohort of pre-school children

Lise Dubois^{1,*}, Anna Farmer², Manon Girard¹, Daniel Burnier¹ and Marion Porcherie¹

¹Department of Epidemiology and Community Medicine, Faculty of Medicine, Institute of Population Health, University of Ottawa, 1 Stewart Street, Office 303, Ottawa, Ontario K1N 6N5, Canada; ²Department of Agricultural, Food and Nutritional Science and the Centre for Health Promotion Studies, University of Alberta, Edmonton, Alberta, Canada

Submitted 16 March 2010; Accepted 8 December 2010; First published online 10 February 2011

Abstract

Objective: To examine: (i) children's food intake and adherence to both Canada's Food Guide for Healthy Eating and Dietary Reference Intakes; and (ii) the social and demographic factors related to children's food intake.

Design: A cross-sectional study.

Setting: Data were obtained through the Quebec Longitudinal Study of Child Development 1998–2010, a representative sample (n 2103) of children born in 1998 in the province of Quebec, Canada. Information on energy, macronutrient and food consumption was derived from responses to a 24 h dietary recall interview addressed to children's mothers and day-care staff when the children were 4 years old.

Subjects: A total of 1549 children aged 4 years who participated in a nutritional sub-study.

Results: The mean daily total energy intake was 6360 kJ (1520 kcal) for girls and 6916 kJ (1653 kcal) for boys. For boys and girls alike, energy intake was comprised of approximately 54% carbohydrates, 31% fats and 15% proteins. The mean number of servings consumed from each of the four essential food groups closely approached the dietary recommendations made by Canada's Food Guide for Healthy Eating; however, <2% of the children in the present study actually met the full dietary guidelines. The dietary intake of pre-school children was associated with socio-economic and demographic factors, most notably mother's level of education, mother's immigrant status and sex of the child.

Conclusions: Diet-related disparities associated with socio-economic and demographic factors exist from as early as 4 years of age.

Keywords
Food intake
Nutritional recommendations
Socio-economic factors

Income is only one among many factors affecting the extent and nature of food purchases⁽¹⁾. Social, cultural and family dynamics all exert an influence on the contents of a given household's shopping basket⁽²⁾. Consumer behaviours, tastes and adherence to nutritional recommendations are socially shaped, and are an object of social distinction^(3,4). Disparities in food consumption are also observed in childhood^(5,6). National nutrition surveys such as the Continuing Survey of Food Intakes by Individuals (CSFII)⁽⁷⁾ and the National Health and Nutrition Examination Survey (NHANES)⁽⁸⁾ have shown that children and adolescents are consuming products high in sugar and fat, and low in nutrient value^(9,10). It is also reported that children and adolescents rarely meet nutritional recommendations for intakes of fruit, vegetables and whole-grain products^(11,12). In their review of the literature on fruit and vegetable intakes among children

and adolescents aged 6–18 years, Rasmussen *et al.*⁽¹³⁾ reported that gender, age, socio-economic position, preferences, parental intake and home availability/accessibility were the determinants supported by the greatest amount of evidence. Regarding ethnicity, studies showed lower fat intakes in immigrants from South Asian countries in comparison with populations born in the USA^(14,15). Most studies also support the idea that high parental education is positively associated with a higher intake of healthy foods^(5,16,17). However, it is unclear whether these associations also play a role in influencing the dietary intakes of younger children. Kranz and Siega-Riz⁽¹⁸⁾ reported an inverse relationship between mother's education and the intake of added sugars in pre-school children. When analysing the diet of >10 000 children aged 3 years from the Avon Longitudinal Study of Pregnancy and Childhood (ALSPAC), North and Emmett⁽¹⁹⁾

*Corresponding author: Email ldubois@uottawa.ca

showed that children's junk food intake was significantly associated with having a younger, less-educated mother and a mother of lower social class. As only a few studies have examined dietary adequacy in pre-school children⁽²⁰⁾, and even fewer have investigated the socio-economic determinants of healthy eating for this age group, the present study contributes to an area that is not well understood. Hence, the aim of the present paper is to examine the social and demographic factors related to children's food intakes and dietary adherence to both Canada's Food Guide to Healthy Eating (1992–2007) and Dietary Reference Intakes (DRI)⁽²¹⁾ in use at the time of the present study.

Methods

The analyses were performed using data from the Quebec Longitudinal Study of Child Development (QLSCD) 1998–2010, conducted by Santé Québec, a division of the Institut de la Statistique du Québec (ISQ) in Canada^(22,23). The QLSCD was established to examine the role of familial and social factors on children's health, as well as on their cognitive and behavioural development. It followed a representative sample (n 2103) of children born in 1998 in the province of Quebec, Canada. The representative sample was chosen by a random selection of children born throughout the year in each public health geographical area of the province, thus minimizing the effect of seasonality and ensuring geographical representation. Twins and children with major diseases or handicaps at birth were excluded from the study. The children were selected from the master birth register and were first examined at 5 months of age (gestational age adjusted for preterm birth) and subsequently at 1-year intervals.

Of the 2103 infants included in the first cycle of the study, 1944 were still part of the survey in 2002. Of this group, 1549 participated in the nutrition evaluation phase of the study. The age of the children included in the present study varied from 44 to 56 months, with a mean age of 49 (SD 3.12) months. Energy, macronutrient and food consumption levels were derived from a 24 h multiple-pass dietary recall interview administered by trained nutritionists in children's homes⁽²⁴⁾. Caregivers were asked to indicate the foods (e.g. type, quantity and recipes) consumed during the 24 h period preceding the interview. For children attending day care (n 390), trained nutritionists went to their home and also queried day-care kitchen staff regarding the child's food intake (e.g. time, meal and quantity) for the same 24 h period. To assess for inter- and intra-child variability in the consumption of energy and macronutrients (carbohydrates, protein and fats), a second 24 h dietary recall was administered to 50% of the sample. All statistical analyses were conducted using adjusted data with results based on usual food consumption patterns and not solely on a single-day recall. A two-step procedure was used: first, a normalizing transformation; and second, a

linear transformation that provides a distribution of intakes without the intra-individual component^(24,25). The 24 h recalls were administered evenly across all days in the week. Energy and macronutrient consumption levels, along with servings of food for each meal and for each child, were evaluated according to the *Canadian Nutrient File*⁽²⁶⁾ and the US Department of Agriculture (USDA) recipe file⁽²⁷⁾, which calculate usual consumption levels of food in accordance with Canada's Food Guide to Healthy Eating. Dietary data were managed using a validated nutrient analysis software developed by Micro Gesta (version 73.0, Quebec, Canada) specifically for Canadian nutritional studies.

On the basis of the results of a literature review, different factors that may relate to the quality of children's diet were selected to examine their role as potential determinants. Family type (single-parent family or not), household annual income (<\$CAN20 000, \$CAN20 000–39 999, \$CAN40 000–59 999 and \geq \$CAN60 000), income level (sufficient, insufficient and very insufficient*), mother's education (no high-school diploma, high-school diploma, college diploma and university degree), mother's age group (<25, 25–29, 30–34 and \geq 35 years) and immigrant status (immigrant or not) were included in the analyses. Other variables such as mother's BMI (from self-reported weight and height), number of overweight/obese parents and sex of the child were also included. Children's level of physical activity and their energy intake were used as control variables. The degree of children's physical activity was measured through one survey question that asked mothers: 'In your opinion, is your child's level of physical activity less than or more than children of the same age and same sex?' Mothers were asked to respond with one of the following categories: 'much higher', 'slightly higher', 'equal', 'slightly less', 'much less than' and 'do not know'. To examine children's dietary adherence to nutritional recommendations, children's food intakes were compared with both Canada's Food Guide to Healthy Eating (1992–2007) and DRI⁽²¹⁾. In correspondence with current American dietary guidelines†, Canada's Food Guide to Healthy Eating in place at the time of the present study recommended the consumption of 5–10 servings of vegetables and fruit, 5–12 servings of grain products, 2–4 servings of milk and alternatives and 2–3 servings of meat and alternatives per day.

Data were weighted by a factor based on the inverse of the selection probability, the probability of non-response and the post-stratification and attrition rates; this ensured

* Families are classified as having 'sufficient income' when the household income is above the low-income threshold determined by Statistics Canada (2006). When income is between 60% and 90% of the low-income threshold, households are classified as having 'insufficient income'; income levels <60% of the low-income threshold are considered as 'very insufficient'. Statistics Canada (2006), *Low Income Cut-offs for 2005 and Low Income Measures for 2004*, catalogue no. 75F0002MIE, vol. 4.

† Dietary Guidelines for Americans. <http://www.cnpp.usda.gov/Publications/DietaryGuidelines/2005/2005DGPolyDocument.pdf>

that data were longitudinally representative of the infants born in 1998 in the population⁽²⁸⁾. To correct for response bias, the weights provided by ISQ were used. The rates obtained through the present study are therefore comparable to those of other surveys on the same population that use weighted responses, independent of the distribution of the sample⁽²⁴⁾. Preliminary analyses indicated that children participating in the nutrition study (n 1549) were representative of not only infants born in 1998 in the province of Quebec but also of children of the same age (4 years) in the Quebec population. Statistical analyses were based on individuals with no missing values for any of the studied variables. Of the 1549 children, 1522 (98% of the sample) were part of the analyses. The impact of missing data was analysed by conducting with-and-without analyses. Given that missing data did not impact the results, children with missing data were excluded from the analyses.

Statistical analyses were conducted using the SAS statistical software package version 8.2 (SAS Institute Inc., Cary, NC, USA). Pairwise mean comparisons were verified using a one-way ANOVA and Tukey's adjustment of P values. Food servings, in categories, were compared using the χ^2 test of association. Adjusted OR as well as their CI were estimated using logistic regression. In the multivariate regression analyses (Tables 2 and 3), socio-demographic variables were selected from the univariate analyses (Table 1) on the basis of their level of significance in relation to the outcome variables. Significance level was set at 5%.

Results

Table 1 presents the association between different factors and dietary consumption in terms of energy, macro-nutrient and food group servings with respect to recommendations made by Canada's Food Guide to Healthy Eating for the studied population. Children's mean daily total energy intake was 6644 kJ (1588 kcal)/d. For boys and girls alike, energy intake was comprised of approximately 54% carbohydrates, 31% fats and 15% proteins. Boys ($P < 0.0001$), children of immigrant mothers ($P = 0.005$), those living in single-parent families ($P = 0.01$) and those from families with very insufficient income ($P = 0.01$) consumed more energy than did other children. In terms of food group servings, children consumed a mean of 3.5 servings of fruit and vegetables, 4 servings of grain products, 2 servings of milk products and 2 servings of meat and alternatives per day (data not presented). Only 17% of children consumed the recommended ≥ 5 servings of fruit and vegetables or grain products per day. Half (47.7%) of the children consumed the recommended 2 servings of milk products per day, whereas 39% of children consumed the recommended 2 servings of meat and alternatives per day. The consumption

of vegetables and fruit was associated with sex of the child ($P = 0.03$), mother's immigrant status ($P = 0.04$) and education ($P \leq 0.0001$), household annual income ($P = 0.0009$) and mother's BMI ($P = 0.004$), whereas the consumption of grain products was associated with sex of the child ($P \leq 0.0001$), mother's immigrant status ($P \leq 0.0001$), single parenting ($P = 0.01$) and family income ($P = 0.02$). The consumption of milk products was associated with family income ($P = 0.01$) and the consumption of meat and alternatives was associated with sex of the child ($P \leq 0.0001$) and single parenting ($P = 0.02$).

The adjusted means for energy intake and macro-nutrient consumption are presented in Table 2. Boys consumed more energy than did girls ($P < 0.0001$), and children of immigrant mothers consumed more energy than did children of non-immigrant mothers ($P < 0.0001$), independently of children's level of physical activity. The proportion of fat consumed was higher in children from non-immigrant mothers ($P = 0.04$) and in children from mothers with no high-school diploma ($P < 0.0001$). The proportion of energy intake derived from protein was higher in children of immigrant mothers ($P = 0.01$) and in children of more highly educated mothers ($P < 0.0001$).

The adjusted OR for consuming ≥ 5 servings of vegetables and fruit per day was 2.2 in children of non-immigrant mothers, in comparison with children of immigrant mothers ($P = 0.002$), and was as high as 8.5 in children of mothers with a university diploma in comparison with children of mothers with no high-school diploma ($P < 0.0001$; Table 3). The adjusted OR for consuming ≥ 5 servings of grain products per day was 5.0 in boys, in comparison with girls ($P < 0.0001$), and 0.4 in children of non-immigrant mothers, in comparison with children of immigrant mothers ($P < 0.0001$). The adjusted OR of consuming ≥ 2 servings of milk products per day was 0.7 in boys in comparison with girls ($P = 0.003$), and 1.5 in children of mothers with a university diploma, in comparison with children of mothers with no high-school diploma ($P = 0.03$). The adjusted OR of consuming ≥ 2 servings of milk products per day in children living in households with insufficient or very insufficient income was 0.6, in comparison with children living in households with sufficient income ($P = 0.0009$). The adjusted OR for consuming ≥ 2 servings of meat and alternatives per day was 1.9 for boys, in comparison with girls ($P < 0.0001$).

Discussion

The present study found that children consumed macro-nutrients in the proportions recommended by Canadian dietary guidelines, which suggest that 45–65% of energy intake should be derived from carbohydrates, 25–35% from fats and 10–30% from proteins⁽²⁹⁾. With regard to daily total energy intake, it is difficult to determine whether the levels reported in the present study are well adapted

Table 1 Mean energy and macronutrient intakes and the prevalence of children attaining the dietary recommendations for four food groups by different characteristics in 4-year-old children from the province of Quebec (from 24 h dietary recall)

Characteristic	Energy		Carbohydrates		Total fats		Proteins		Fruit and vegetables	Grain products	Milk products	Meat and alternatives
	kcal	SD	%E	SD	%E	SD	%E	SD	≥5 servings	≥5 servings	≥2 servings	≥2 servings
Total	1588	7	53.8	0.1	31.4	0.1	14.5	0.1	16.6	16.8	47.7	39.1
Child's sex												
Girl	1520*	10	53.8	0.1	31.4	0.1	14.5	0.1	14.5*	7.0*	46.7	28.2*
Boy	1653	9	53.8	0.1	31.4	0.1	14.5	0.1	18.6	26.3	48.6	49.6
Mother's age (years)												
<25	1613	26	53.3	0.3	31.9	0.2	14.4	0.2	13.9	20.7	47.9	43.5
25–29	1612	14	54.0	0.2	31.3	0.1	14.3	0.1	14.1	16.2	44.5	40.2
30–34	1575	12	53.8	0.2	31.4	0.1	14.5	0.1	19.9	15.7	46.4	37.6
≥35	1579	11	53.7	0.1	31.3	0.1	14.7	0.1	16.1	17.2	50.8	39.5
Mother's immigrant status												
Not immigrant	1579*	7	53.8	0.1	31.5*	0.1	14.5*	0.1	17.5*	14.4*	47.0	38.9
Immigrant	1635	19	54.0	0.2	30.9	0.2	14.8	0.1	11.8	31.1	53.1	41.4
Mother's education												
No high-school diploma	1622	17	53.7	0.2	31.8*	0.2	14.0*	0.1	7.7*	17.1	41.8	37.7
High-school diploma	1575	15	53.9	0.2	31.4	0.1	14.4	0.1	11.4	15.4	46.3	39.0
College diploma	1581	12	53.7	0.2	31.4	0.1	14.6	0.1	14.1	17.9	50.0	39.4
University diploma	1586	13	53.8	0.2	31.2	0.1	14.8	0.1	29.7	16.1	49.3	40.6
Single parenting												
No	1581*	7	53.7	0.1	31.4	0.1	14.5	0.1	16.2	16.0*	48.0	38.0*
Yes	1628	18	54.1	0.2	31.1	0.2	14.5	0.1	19.1	22.5	45.9	45.7
Annual income (\$CAN)												
<20 000	1596	21	53.9	0.3	31.2	0.2	14.5*	0.1	13.3*	14.1	39.4	47.7
20 000–39 999	1604	15	53.8	0.2	31.5	0.1	14.3	0.1	13.1	20.9	46.5	36.9
40 000–59 999	1600	13	53.7	0.2	31.6	0.1	14.3	0.1	14.0	16.0	47.8	39.1
≥60 000	1568	11	53.8	0.1	31.2	0.1	14.7	0.1	21.6	16.1	50.7	38.8
Income level												
Sufficient	1579*	8	53.7	0.1	31.4	0.1	14.5	0.1	17.1	16.0*	49.6*	38.0
Insufficient	1613	19	53.9	0.3	31.3	0.2	14.4	0.1	17.3	24.0	38.5	41.9
Very insufficient	1650	26	53.8	0.4	31.3	0.2	14.4	0.2	11.9	14.0	43.3	50.9
Mother's BMI (kg/m ²)												
<18.5	1593	27	54.2	0.4	31.0	0.2	14.5	0.2	22.9*	17.4	44.8	40.7
18.5–24.9	1587	9	53.7	0.1	31.4	0.1	14.6	0.1	17.5	15.2	48.3	38.1
25.0–29.9	1586	16	54.0	0.2	31.3	0.1	14.5	0.1	16.9	18.7	45.4	39.4
≥30.0	1586	23	53.9	0.3	31.5	0.2	14.1	0.2	6.6	22.5	47.5	40.8
Number of overweight/obese parents												
0	1583	11	53.9	0.1	31.3	0.1	14.5	0.1	19.1	16.0	47.4	37.4
1	1585	10	53.8	0.1	31.4	0.1	14.5	0.1	15.9	16.3	48.0	38.4
2	1596	18	53.8	0.2	31.5	0.2	14.4	0.1	13.1	18.8	45.6	42.8

%E, percentage of energy.

1 kcal = 4.184 kJ.

*Statistically significant association between the characteristic and the dependent variable (P value ≤ 0.05).

Table 2 Adjusted mean energy and macronutrient intakes by different characteristics

Characteristic	Energy†		Carbohydrates‡		Total fats‡		Proteins‡	
	kcal	SD	%E	SD	%E	SD	%E	SD
Child's sex								
Girl	1562*	39	53.4	0.5	31.5	0.4	14.7	0.3
Boy	1690	38	53.4	0.5	31.5	0.4	14.6	0.3
Mother's immigrant status								
Not immigrant	1603*	38	53.4	0.5	31.7*	0.4	14.5*	0.3
Immigrant	1649	41	53.4	0.5	31.3	0.4	14.8	0.3
Mother's education								
No high-school diploma	1656	41	53.4	0.6	31.9*	0.4	14.2*	0.3
High-school diploma	1605	40	53.5	0.6	31.4	0.4	14.6	0.3
College diploma	1616	39	53.3	0.5	31.4	0.4	14.8	0.3
University diploma	1627	40	53.5	0.5	31.2	0.4	15.0	0.3
Single parenting								
No	1612	38	53.3	0.5	31.6	0.4	14.6	0.3
Yes	1639	41	53.5	0.6	31.3	0.4	14.7	0.3
Income level								
Sufficient	1606	39	53.3	0.5	31.6	0.4	14.7	0.3
Insufficient	1610	40	53.4	0.5	31.4	0.4	14.6	0.3
Very insufficient	1661	45	53.4	0.6	31.4	0.4	14.6	0.3

%E, percentage of energy.

1 kcal = 4.184 kJ.

*Statistically significant association between the characteristic and the dependent variable (P value ≤ 0.05).

†Adjusted for child's sex, mother's immigrant status, mother's education, single parenting, income level and physical activity level.

‡Adjusted for child's sex, mother's immigrant status, mother's education, single parenting, income level, and physical activity level, and carbohydrates, lipids and proteins as proportion of energy.

Table 3 Adjusted OR for attaining dietary recommendations for food group servings at 4 years of age by different characteristics

Characteristic	Fruit and vegetable†		Grain product†		Milk product†		Meat and alternativest	
	≥ 5 servings		≥ 5 servings		≥ 2 servings		≥ 2 servings	
	Adjusted OR	95% CI	Adjusted OR	95% CI	Adjusted OR	95% CI	Adjusted OR	95% CI
Child's sex								
Girl	1.0	Ref.	1.0	Ref.	1.0	Ref.	1.0	Ref.
Boy	1.1	0.8, 1.5	5.0*	3.3, 7.5	0.7*	0.5, 0.9	1.9*	1.5, 2.4
Mother's immigrant status								
Not immigrant	2.2*	1.3, 3.7	0.4*	0.2, 0.6	0.7	0.5, 1.1	1.2	0.9, 1.7
Immigrant	1.0	Ref.	1.0	Ref.	1.0	Ref.	1.0	Ref.
Mother's education								
No high-school diploma	1.0	Ref.	1.0	Ref.	1.0	Ref.	1.0	Ref.
High-school diploma	2.0*	1.0, 3.7	1.1	0.6, 1.9	1.6*	1.1, 2.4	1.3	0.9, 1.9
College diploma	2.8*	1.5, 5.0	1.4	0.8, 2.4	1.7*	1.2, 2.5	1.3	0.9, 1.9
University diploma	8.5*	4.6, 15.4	1.1	0.6, 2.0	1.5*	1.0, 2.3	1.5	1.0, 2.2
Single parenting								
No	1.0	Ref.	1.0	Ref.	1.0	Ref.	1.0	Ref.
Yes	1.3	0.8, 1.9	1.4	0.9, 2.3	1.0	0.7, 1.4	1.3	0.9, 1.8
Income level								
Sufficient	1.0	Ref.	1.0	Ref.	1.0	Ref.	1.0	Ref.
Insufficient/very insufficient	0.8	0.6, 1.3	0.8	0.5, 1.4	0.6*	0.4, 0.8	1.3	0.9, 1.8

Ref., reference category.

*Statistically significant OR and 95% CI (P value ≤ 0.05).

†Adjusted for child's sex, mother's immigrant status, mother's education, single parenting, income level, energy and physical activity level.

to this population; energy intake is dependent upon children's level of physical activity, and this was documented through parental report and not directly measured⁽²³⁾. However, the observed levels of energy intake concur with values observed through the NHANES. Across the years 1999–2000, NHANES reported that children aged 3–5 years consumed a mean daily energy intake of 6786 kJ (1622 kcal)⁽³⁰⁾. American figures (NHANES) and those reported in the present study are clearly higher

than what is advised by the Nordic Nutrition Recommendations for children aged 2–5 years, which stipulate a daily energy intake of 5293 kJ (1265 kcal)⁽³¹⁾. Using various dietary methodologies and sample sizes, a summary of European studies reported by Lambert *et al.*⁽³²⁾ allows for other comparisons. For instance, energy intakes for boys aged 4–6 years are reported to range from 5293 kJ (1265 kcal)/d in the UK to 7694 kJ (1839 kcal)/d in Austria. Corresponding intakes for girls vary from 5096 kJ (1218 kcal)/d to 9590 kJ

(2292 kcal)/d across these same countries. For 4–6-year-old boys, the proportion of energy intakes derived from carbohydrates is reported to vary from 44% in Greece to 56.8% in Russia. As a percentage of total energy, the fat intakes of 4–6-year-old girls are also reported to vary from 35.5% in the UK to 41% in Greece, whereas protein intakes vary from 11.7% in Russia to 13.6% in the Netherlands. In the American survey (NHANES), 13.2% of energy intake came from proteins, 55.4% from carbohydrates and 32.9% from fats in children under 6 years of age⁽⁸⁾. The results of the present study suggest that the contribution of protein to children's energy intakes exceeds levels reported in most European studies reviewed by Lambert *et al.*⁽³²⁾. On the contrary, intakes of fat, as a percentage of energy intake, appear to be lower in Quebec than in most European countries. However, total energy intake and the percentage of energy intake from carbohydrates in the present study are comparable to those of most studies.

The mean number of servings of foods consumed from each of the four essential food groups by children of this cohort closely approached the dietary recommendations made by Canada's Food Guide to Healthy Eating; however, <2% of children actually met the dietary guidelines of all four food groups simultaneously (data not presented). Unlike the findings observed in the present study, a Spanish study of 1112 children aged 6–7 years found that a majority of children maintained the dietary recommendations made by the USDA⁽³³⁾. Compared with findings observed in the USA through a study of children included in the CSFII, the mean number of servings of milk products consumed by children in the present study cohort was equal to the number of servings consumed by children in the CSFII, but was lower for servings of grain products and meat and alternatives⁽³⁴⁾. Compared with results from the ALSPAC study, a cohort of 1026 children aged 1.5 years, the percentage of vegetable/fruit non-consumers (2.5%) between studies is comparable⁽³⁵⁾ (data not presented).

The present study indicates that mother's immigrant status, mother's level of education and sex of the child are especially central socio-economic and demographic factors that relate to children's food intakes. With regard to mother's immigrant status, the findings of the present study concur partially with a study from the Canadian province of Ontario where the immigrant population of adults is reported to consume less fat and more carbohydrates^(36,37). As Pomerleau *et al.*^(36,37) reported, the differences between the diets of children of immigrant mothers and children of non-immigrant mothers may reflect food consumption patterns of different countries of origin, which may also relate to food availability within those countries. For instance, many studies have reported that immigrants from South Asian countries have lower fat intakes than non-immigrants⁽¹⁴⁾. Since immigrants are often in a situation of cultural and dietary transition, various studies have examined the association between

this transition and dietary patterns. For example, Renzaho *et al.*⁽³⁸⁾ recently reported that the maintenance of traditional cultural orientation among African migrant children (3–12-year-olds) to Australia was associated with lower rates of obesity and sedentary behaviours. Given that in the present study cohort only 2.8% of mothers were from European countries whereas 11% originated from non-European countries⁽²⁴⁾, the results suggest that, in this population, immigrants may still be adhering to traditional diets that are high in grains. Children of immigrant mothers are also more likely to be raised in a low-income family compared with children of Canadian-born mothers⁽³⁹⁾, and this may explain why in the present study it was observed that children of immigrant mothers consumed less fruit and vegetables than did children of non-immigrant mothers. Some studies report that lower-income families tend to consume fewer fruit and vegetables and less varied diets^(40,41). However, in the present study, income level was controlled for and the relationship between a mother's immigrant status and children's lower fruit and vegetable intake remained. Thus, it is possible that the dietary choices of new immigrants may be influenced by several other determinants, apart from income level and poverty, including culture, access to information, availability of healthy and acceptable foods and acculturation⁽⁴²⁾.

With regard to the association between mother's level of education and children's food intake, our results are in accordance with findings from the British National Diet and Nutrition Study that analysed the diet of 1675 children aged 1.5–4.0 years; this aforementioned study found that children of mothers with a low level of education and children from families where the head of the household had a manual occupation were less likely to meet the recommended guidelines for macronutrient intakes⁽⁴³⁾. Most studies support a positive association between parental level of education and intakes of healthy foods^(5,16,17). Since recent studies⁽⁴⁾ have reported that different socio-economic groups are largely familiar with nutritional recommendations, it may be hypothesized that nutritional recommendations (such as other medical recommendations regarding child care, smoking, etc.) may be valued differently in lower socio-economic groups. Furthermore, as nutritional recommendations are usually encouraged and advocated by medical professionals and individuals from higher socio-economic groups, such recommendations may be perceived by individuals from lower socio-economic groups as an attempt to impose a moral constraint or 'an attempt to impose a control on their bodies or the ones of their children'⁽⁴⁾, as such individuals may be more concerned with meeting daily needs or maintaining other values and pleasures of life. Further research is needed to examine this hypothesis.

Finally, the findings of the present study on the association between the sex of a child and food intake contrast with findings from the ALSPAC study, which found

that, among 7-year-old children, girls ate more fruit and vegetables in comparison with boys⁽⁴⁴⁾. However, as was observed in the present study, being a boy was associated with a higher intake of grain products. It is possible that, as was found in the ALSPAC study, boys may be consuming greater quantities of breakfast cereals in comparison with girls, who appear to be more likely to skip breakfast than boys⁽⁴⁵⁾. However, this hypothesis remains to be tested. With regard to fruit and vegetable intakes, in 1998 the CSFII survey found that boys consumed more of all foods except fruit and vegetables⁽⁷⁾. However, using data from the 1999–2002 NHANES, Lorson *et al.*⁽⁴⁶⁾ recently reported that among children aged 2–18 years of age boys consumed significantly more vegetables than did girls. As boys also consumed significantly more French fries than did girls (the leading source of vegetables in the study reported by Lorson *et al.*⁽⁴⁶⁾), it should not be automatically assumed that boys consumed more dark green or orange vegetables than did girls*. At this pre-school age, children's preferences and differences between boys and girls are certainly influenced by the food environment that the parent provides. In a study on 3–4-year-old children, Johnson and Birch⁽⁴⁷⁾ reported differences in how boys and girls are parented regarding food and eating practices. Their results showed that 'restrained' mothers attempted to restrain their daughters and not their sons. These results indicate that differential pressure for thinness between boys and girls begins during the pre-school period. These differences in parental child-feeding practices may also explain why boys were found to consume more energy than were girls in the present study. Analyses conducted by Dubois and Girard⁽⁴⁸⁾ using data from the QLSCD showed that a higher proportion of mothers overestimated weights for boys than for girls. The authors hypothesized that mothers of young children may be influenced by social desirability and have different standards of what is acceptable in weight for boys and girls. It is also possible that boys' energy needs are higher than that of girls because of larger body height and weight. In fact, Desrosiers *et al.*⁽²⁴⁾, who also used data from the QLSCD, reported sex differences in mean weight and height even at the age of 4 years: 17.2 kg for boys *v.* 16.6 kg for girls; 104 cm for boys *v.* 102.8 for girls. Finally, the fact that children's level of physical activity was incorporated among the control variables is limited by the reported rather than measured nature of this variable.

Strengths, limitations and conclusion

The results of the present study should be considered in the context of its strengths and limitations. The large

representative population-based sample and the high response rate (85%) obtained are the most important strengths of the present study. Although the 24 h recall method is considered to be one of the best measures used to estimate dietary intake, it is known that this method may be biased by a high proportion of under-reporters⁽⁴⁹⁾. For instance, there is an increased tendency among obese parents to under-report children's dietary intakes in comparison with non-obese parents⁽⁵⁰⁾. Some studies also report that snack foods are commonly under-reported⁽⁵¹⁾. To avoid these biases, O'Connor *et al.*⁽⁵²⁾ recommend that children aged 6–9 years complete three 24 h recall interviews to minimize errors associated with under-reporting. Although we did not reach this recommended objective, energy, macronutrient and food consumption patterns were derived from a 24 h dietary recall interview conducted by trained nutritionists and a second 24 h dietary recall was administered to 50% of the sample to assess for inter- and intra-child variability in the consumption of energy and macronutrients. The absence of information on parental food intake is a noteworthy limitation as it has been shown that parental fruit and vegetable intake is a strong predictor of children's fruit and vegetable intake⁽⁵³⁾. However, given that diet quality is tributary to socio-economic factors such as those measured in the present study⁽⁵⁴⁾, it is assumed that the lack of this information has a minimal impact on the results. Another limitation to the study results is the absence of information on children's weight and height.

Overall, the present study revealed that even for pre-school children dietary intakes were associated with socio-economic and demographic factors, most notably with mother's level of education, immigrant status and sex of the child. Children of mothers with no high-school diploma consumed fewer proteins, more fats (by percentage of total energy intake) and were less likely to consume the recommended servings of fruit and vegetables in comparison with children of mothers with a higher level of education. Children of immigrant mothers consumed significantly more energy but less fruit and vegetables than did children of non-immigrant mothers. The present study also revealed that the odds for consuming the recommended servings of grain products and meat and alternatives were higher in boys than in girls, whereas for milk products the odds were higher in girls than in boys. Reducing diet-related disparities and promoting healthy eating involve a complex interaction of factors. When efforts to improve the nutrition and health of young children are planned, these diet-related disparities need to be taken into account.

Acknowledgements

The research work of L.D. was supported by the Canada Research Chair Program. The present study has been

* For vegetables classification, see MyPyramid on the USDA website: www.mypyramid.gov

partly financed by the Canadian Institute of Health Information, Population Health Initiative and by the Canadian Institute of Health Research. Analyses were performed using data from the Longitudinal Study of Child Development in Québec (LSCDQ) 1998–2010, conducted by Santé Québec, a division of the Institut de la Statistique du Québec, and funded by the Ministry of Health and Social Services of Québec. No conflict of interest is declared. L.D. is the director of the study (questionnaire development, data analysis, interpretation of the data, etc.); M.G. planned and performed the statistical data analysis; A.F., D.B. and M.P. contributed to literature review, content analysis and writing of the manuscript. All authors have seen and approved the manuscript and have met all requirements for authorship.

References

- Halbwachs M (1938) *Esquisse d'une Psychologie des Classes Sociales*. Paris: Presses Universitaires de France.
- Régnier F (2009) Obésité, goûts et consommation. Intégration des normes d'alimentation et appartenance sociale. *Rev Fr Sociol* **50**, 747–773.
- Bourdieu P (1984) *Distinction: A Social Critique of the Judgement of Taste*. Cambridge, MA: Harvard University Press.
- Régnier F & Masullo A (2008) *Une Affaire de Goût? Réception et Mise en Pratique des Recommandations Nutritionnelles*. Paris: INRA.
- Roos G, Johansson L, Kasmel A *et al.* (2001) Disparities in vegetable and fruit consumption: European cases from the north to the south. *Public Health Nutr* **4**, 35–43.
- Taylor JP, Evers S & McKenna M (2005) Determinants of healthy eating in children and youth. *Can J Public Health/Rev Can Sante Publique* **96**, Suppl. 3, S20–S26.
- Wilkinson-Enns C, Mickle SJ & Goldman JD (2002) Trends in food and nutrient intakes by children in the United States. *Fam Econ Nutr Rev* **14**, 56–68.
- National Centre for Health Statistics (2003) Dietary intake of ten key nutrients for public health, United States: 1999–2000. *Adv Data* **334**, 1–4.
- Kant AK & Graubard BI (2003) Predictors of reported consumption of low-nutrient-density foods in a 24-h recall by 8–16 year old US children and adolescents. *Appetite* **41**, 175–180.
- Subar AF, Krebs-Smith SM, Cook A *et al.* (1998) Dietary sources of nutrients among US children, 1989–1991. *Pediatrics* **102**, 913–923.
- Troiano RP, Briefel RR, Carroll MD *et al.* (2000) Energy and fat intakes of children and adolescents in the United States: data from the National Health and Nutrition Examination Surveys. *Am J Clin Nutr* **72**, Suppl. 5, 1343S–1353S.
- Skinner JD, Carruth BR, Houck KS *et al.* (1999) Longitudinal study of nutrient and food intakes of white preschool children aged 24 to 60 months. *J Am Diet Assoc* **99**, 1514–1521.
- Rasmussen M, Krolner R, Klepp K-I *et al.* (2006) Determinants of fruit and vegetable consumption among children and adolescents: a review of the literature. Part I: quantitative studies. *Int J Behav Nutr Phys Act* **3**, 22.
- Kim KK, Yu ES, Liu WT *et al.* (1993) Nutritional status of Chinese-American, Korean-American, and Japanese-American elderly. *J Am Diet Assoc* **93**, 1416–1422.
- Xie B, Gilliland FD, Li YF *et al.* (2003) Effects of ethnicity, family income, and education on dietary intake among adolescents. *Prev Med* **36**, 30–40.
- Sausenthaler S, Kompauer I, Mielck A *et al.* (2007) Impact of parental education and income inequality on children's food intake. *Public Health Nutr* **10**, 24–33.
- Serra-Majem L, Ribas L, Perez-Rodrigo C *et al.* (2002) Determinants of nutrient intake among children and adolescents: results from the enKid study. *Ann Nutr Metab* **46**, 31–38.
- Kranz S & Siega-Riz AM (2002) Sociodemographic determinants of added sugar intake in preschoolers 2 to 5 years old. *J Pediatr* **140**, 667–672.
- North K & Emmett P (2000) Multivariate analysis of diet among three-year-old children and associations with socio-demographic characteristics. *Eur J Clin Nutr* **54**, 73–80.
- Nicklaus S, Boggio V *et al.* (2004) A prospective study of food preferences in childhood. *Food Qual Prefer* **15**, 805–818.
- Institute of Medicine (1998) *Dietary Reference Intakes: A Risk Assessment Model for Establishing Upper Intake Levels for Nutrients*. Washington, DC: National Academy Press.
- Dubois L, Bédard B, Girard M *et al.* (2000) *Etude Longitudinale du Développement des Enfants du Québec (ELDEQ 1998–2002). Les Nourrissons de 5 Mois*. Québec: Institut de la Statistique du Québec.
- Dubois L & Girard M (2002) *L'alimentation des Enfants d'Âge Préscolaire. Etude Longitudinale du Développement des Enfants du Québec (ELDEQ 1998–2002)*. Québec: Institut de la Statistique du Québec.
- Desrosiers H, Dubois L, Bédard B *et al.* (2005) *Enquête de Nutrition Auprès des Enfants Québécois de 4 Ans*. Québec: Institut de la Statistique du Québec.
- Liu K, Stampler J, Dyer A *et al.* (1978) Statistical methods to assess and minimize role of intra-individual variability in obscuring relationship between dietary lipids and serum cholesterol. *J Chron Dis* **31**, 399–418.
- Health Canada (2001) *Canadian Nutrient File*. Ottawa: Health Canada.
- US Department of Agriculture (2004) *Food and Nutrient Database for Dietary Studies, 1.0. 2004*. Beltsville, MD: Agricultural Research Service, Food Surveys Research Group.
- Cox B & Cohen S (1985) *Methodological Issues for Health Care Surveys*. New York: Marcel Dekker.
- Institute of Medicine (2002) *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein and Amino Acids*. Washington, DC: National Academy Press.
- Briefel RR & Johnson CL (2004) Secular trends in dietary intake in the United States. *Ann Rev Nutr* **24**, 401–431.
- Nordic Council of Ministers (2004) *Nordic Nutrition Recommendations 2004. Integrating Nutrition and Physical Activity*. Copenhagen: NCM.
- Lambert J, Agostoni C, Elmadfa I *et al.* (2004) Dietary intake and nutritional status of children and adolescents in Europe. *Br J Nutr* **92**, Suppl. 2, S147–S211.
- Royo-Bordonada MA, Gorgojo L, Martin-Moreno JM *et al.* (2003) Spanish children's diet: compliance with nutrient and food intake guidelines. *Eur J Clin Nutr* **57**, 930–939.
- Munoz KA, Krebs-Smith SM, Ballard-Barbash R *et al.* (1997) Food intakes of US children and adolescents compared with recommendations. *Pediatrics* **100**, 323–329.
- Cowen I & Emmett P (2000) Diet in a group of 18-month-old children in South West England, and comparison with the results of a national survey. *J Hum Nutr Diet* **13**, 87–100.
- Pomerleau J, Ostbye T & Bright-See E (1998) Place of birth and dietary intake in Ontario I. Energy, fat, cholesterol, carbohydrate, fiber, and alcohol. *Prev Med* **27**, 32–40.
- Pomerleau J, Ostbye T & Bright-See E (1998) Place of birth and dietary intake in Ontario II. Protein and selected micronutrients. *Prev Med* **27**, 41–49.
- Renzaho AMN, Swinburn B & Burns C (2008) Maintenance of traditional cultural orientation is associated with lower rates of obesity and sedentary behaviours among African migrant children to Australia. *Int J Obes (Lond)* **32**, 594–600.

39. Statistics Canada (2003) *The Rise in Low-Income Rates Among Immigrants in Canada*. Ottawa: Statistics Canada.
40. Kirkpatrick S & Tarasuk V (2003) The relationship between low income and household food expenditure patterns in Canada. *Public Health Nutr* **6**, 89–97.
41. Rogers I & Emmett P (2003) The effect of maternal smoking status, educational level and age on food and nutrient intakes in preschool children: results from the Avon Longitudinal Study of Parents and Children. *Eur J Clin Nutr* **57**, 854–864.
42. Hyman I, Guruge S, Makarchuk MJ *et al.* (2002) Promotion of healthy eating among new immigrant women in Ontario. *Can J Diet Pract Res* **63**, 125–129.
43. Watt RG, Dykes J & Sheiham A (2001) Socio-economic determinants of selected dietary indicators in British pre-school children. *Public Health Nutr* **4**, 1229–1233.
44. Glynn L, Emmett P & Rogers I (2005) Food and nutrient intakes of a population sample of 7-year-old children in the south-west of England in 1999/2000 – what difference does gender make? *J Hum Nutr Diet* **18**, 7–19.
45. Rampersaud GC, Pereira MA, Girard BL *et al.* (2005) Review – breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. *J Am Diet Assoc* **105**, 743–760.
46. Lorson BA, Melgar-Quinonez HR & Taylor CA (2009) Correlates of fruit and vegetable intakes in US children. *J Am Diet Assoc* **109**, 474–478.
47. Johnson SL & Birch LL (1994) Parents and childrens adiposity and eating style. *Pediatrics* **94**, 653–661.
48. Dubois L & Girard M (2007) Accuracy of maternal reports of pre-schoolers' weights and heights as estimates of BMI values. *Int J Epidemiol* **36**, 132–138.
49. Black AE, Goldberg GR, Jebb SA *et al.* (1991) Critical evaluation of energy-intake data using fundamental principles of energy physiology. 2. Evaluating the results of published surveys. *Eur J Clin Nutr* **45**, 583–599.
50. Heitmann BL & Lissner L (2005) Can adverse effects of dietary fat intake be overestimated as a consequence of dietary fat underreporting? *Public Health Nutr* **8**, 1322–1327.
51. Krebs-Smith SM, Graubard BI, Kahle LL *et al.* (2000) Low energy reporters vs others: a comparison of reported food intakes. *Eur J Clin Nutr* **54**, 281–287.
52. O'Connor J, Ball EJ, Steinbeck KS *et al.* (2001) Comparison of total energy expenditure and energy intake in children aged 6–9 y. *Am J Clin Nutr* **74**, 643–649.
53. Cooke LJ, Wardle J, Gibson EL *et al.* (2004) Demographic, familial and trait predictors of fruit and vegetable consumption by pre-school children. *Public Health Nutr* **7**, 295–302.
54. Northstone K & Emmett P (2005) Multivariate analysis of diet in children at four and seven years of age and associations with socio-demographic characteristics. *Eur J Clin Nutr* **59**, 751–760.